arXiv:1505.00367v1 [math.CV] 2 May 2015

EXTENSION OF L^2 HOLOMORPHIC FUNCTIONS

LUCA BARACCO

Abstract. The purpose of this note is to show that the $\bar\partial$-estimate which is needed in the Ohsawa-Takegoshi Extension Theorem [6] is a direct consequence of the Hörmander-Kohn-Morrey weigthed inequality. In this inequality, the Donnelly-Fefferman argument is not required and a single 1-parameter family of non-singular weights is used. This paper is the furtherst step of a great deal of work devoted to the simplification of the original proof of Ohsawa-Takegoshi Theorem; among other papers on the subject, we mention [1] and [8] which are based on "twisted" basic estimates and, in recent time, [3] and [9].

MSC: 32F10, 32F20, 32N15, 32T25

Through an approximation argument, the extension theorem relies on Theorem 1 below.

Theorem 1. *Let $D\subset\subset\mathbb{C}^n$ be a bounded smooth pseudoconvex domain with diameter ≤ 1, ψ a plurisubharmonic function on D, α a $\bar\partial$-closed form in $L^2_\psi(D)$ of degree ≥ 1 such that $\alpha_J=0$ for $1\notin J$ and $\operatorname{supp}\alpha\subset\{z:|z_1|<\delta\}$. Then there is a solution $u=u_\delta\in L^2_\psi$ to the problem*

$$\begin{cases}\bar\partial u=\alpha,\\ \|z_1u\|_\psi\le c\delta\|\alpha\|_\psi, \quad \textit{for } c \textit{ independent of } \delta,\ \psi \textit{ and } D.\end{cases} \tag{1}$$

Proof. We first assume $\psi=0$. We set

$$\varphi(=\varphi^\delta)=-\log(-\log(|z_1|^2+\delta^2)),\qquad a(=a^\delta)=-\log(|z_1|^2+\delta^2); \tag{2}$$

they are related by $a=e^{-\varphi}$. Their key properties are

$$\varphi^\delta_1=\frac{-\bar z_1}{\log(|z_1|^2+\delta^2)(|z_1|^2+\delta^2)},\quad \varphi^\delta_{1\bar1}=\frac{|z_1|^2-\delta^2\log(|z_1|^2+\delta^2)}{\log^2(|z_1|^2+\delta^2)(|z_1|^2+\delta^2))^2}. \tag{3}$$

This readily implies

$$-a_{1\bar1}\ge 0,\quad -a_{1\bar1}\underset{\sim}{>}\frac{1}{\delta^2}\text{ for }|z_1|<\delta. \tag{4}$$

The contraction of the gradient with a multivector v is defined by $\partial a\,\llcorner\, v=(a_i)\,\llcorner\, v=\sum_i a_iv_{iK}$ and the action of the Levi form is $\partial\bar\partial a(v,v)=(a_{i\bar j})(v,v)=\sideset{}{'}\sum\limits_{|K|=k-1}\sum\limits_{ij=1,\dots,N}a_{i\bar j}v_{iK}\bar v_{jK}$.

From the identity $\bar\partial^*_\varphi=\bar\partial^*+\partial\varphi\,\llcorner$, we get the estimate

$$\|\bar\partial^*_\varphi v\|^2_\varphi\le\|\bar\partial^*v\|^2_\varphi+\|\partial\varphi\,\llcorner\, v\|^2_\varphi+2\Big|\int_D e^{-\varphi}(\partial\varphi\,\llcorner\, v)\cdot(\bar\partial^*v)dV\Big|. \tag{5}$$

We also have

$$\begin{aligned}2\Big|\int_D e^{-\varphi}\partial\varphi\llcorner v\,\cdot\,\bar\partial^* v\,dV\Big| &\underset{(2)}{=} 2\Big|\int_D \partial a\llcorner v\cdot\bar\partial^* v dV\Big|\\ &\underset{\text{Cauchy-Schwarz}}{\le} \|v\|_0^2+\||\partial a|\bar\partial^* v\|_0^2\\ &\underset{\text{Pseudoconvexity}}{\lesssim} c(\|\bar\partial v\|_0^2+\|\bar\partial^* v\|_0^2+\||\partial a|\bar\partial^* v\|_0^2),\end{aligned}\tag{6}$$

where c only depends on the diameter of D. Recall that $a=e^{-\varphi}$ and $\|\cdot\|_\varphi=\|a^{\frac12}\cdot\|_0$. We may then conclude

$$\begin{aligned}\int_D -(a_{i\bar j})(v,v)dV &= \int e^{-\varphi}\varphi_{i\bar j}(v,v)dV-\|\partial\varphi\llcorner u\|_\varphi^2\\ &\underset{\text{basic}}{\le} \|\bar\partial v\|_\varphi^2+\|\bar\partial_\varphi^* v\|_\varphi^2-\|\partial\varphi\llcorner u\|_\varphi^2\\ &\underset{(5)}{\le} \|\bar\partial v\|_\varphi^2+\|\bar\partial^* v\|_\varphi^2+2\Big|\int_D e^{-\varphi}\partial\varphi\llcorner v\,\cdot\,\bar\partial^* v\,dV\Big|\\ &\le c(\|(1+a^{\frac12})\bar\partial v\|_0^2+\|(1+a^{\frac12}+|\partial a|)\bar\partial^* v\|_0^2).\end{aligned}\tag{7}$$

With (7) in our hands, we define a functional in L^2 by putting

$$(1+a^{\frac12}+|\partial a|)\bar\partial^* v\mapsto (v,\alpha)_0\quad\text{for } v\in D_{\bar\partial^*}.\tag{8}$$

If $v\in(\ker\bar\partial)^\perp$, we have $(v,\alpha)_0=0$. If, instead, $v\in\ker\bar\partial$, then

$$\begin{aligned}|(v,\alpha)_0| &= \Big|\int_D(-a_{i\bar j})(v,(-a_{1\bar 1})^{-1}\alpha)dV\Big|\\ &\underset{\text{Cauchy-Schwarz}}{\le} \left(\int_D(-a_{i\bar j})(v,v)dV\right)^{\frac12}\|(-a_{1\bar1})^{-\frac12}\alpha\|_0\\ &\underset{(4)}{\le} \left(\int_D(-a_{i\bar j})(v,v)dV\right)^{\frac12}(\delta\|\alpha\|_0)\\ &\underset{(7)}{\lesssim} \|(1+a^{\frac12}+|\partial a|)\bar\partial^* v\|\,\delta\|\alpha\|_0,\end{aligned}\tag{9}$$

where, in order to apply Cauchy-Schwarz, we have used that $(-a_{i\bar j})\ge 0$. Thus, by Riesz Representation Theorem, there exists $w\in L^2$ of norm $\|w\|_0\lesssim\delta\|\alpha\|_0$ which represents (8). Setting $u:=(1+a^{\frac12}+|\partial a|)w$, we get $\bar\partial u=\alpha$; also, from the trivial inequality $(1+a^{\frac12}+|\partial a|)\le|z_1|^{-1}$, we conclude $\|z_1u\|_0\lesssim\delta\|\alpha\|_0$. (Note that we replace $(1+a^{\frac12}+|\partial a|)$ by $|z_1|^{-1}$ in the conclusion but not in the proof, especially in (9), because $|z_1|^{-1}$ is not in L^2.)

When ψ is smooth in $\bar{D}$, the proof above can be repeated verbatim, with the weight φ replaced by $\varphi+\psi$ in the basic estimate (7) and with the Levi form (ψ_{ij}) dropped down because of its positivity. For a general plurisubharmonic ψ, we take a smooth approximation of ψ from above and an exhaustion of D from inside. A weak limit of solutions on these subdomains yields the solution on D. For this approximation we need that $L^2_\psi \subset L^2_{\text{loc}}$. But this follows from the local boundedness of ψ from above which is in turn a consequence of its upper semicontinuity.

□

REFERENCES

[1] **B. Berndtsson**—The extension theorem of Ohsawa-Takegoshi and the theorem of Donnelly-Fefferman, *Ann. Inst. Fourier* **46** n. 4 (1996), 1083–1094

[2] **Z. Blocki**—Cauchy-Riemann meet Monge-Ampère, *Bull. Math. Sci.* **4** (2014), 433–480

[3] **B.Y. Chen**—A simple proof of the Ohsawa-Takegoshi extension theorem, arxiv : 11052430v1

[4] **L. Manivel**—Un théorème de prolongement L^2 des sections holomorphes d'un fibré hermitien, *Math. Z.* **212** n. 1 (1993), 107–122

[5] **J.D. McNeal and D. Varolin**—L^2 extension of $\bar\partial$-closed forms from a hypersurface, arXiv:1502.08054v1

[6] **T. Ohsawa**—On the extension of L^2 holomorphic functions II, *Publ. RIMS* **24** (1988), 265–275

[7] **T. Ohsawa and K. Takegoshi**—On the extension of L^2, holomorphic functions, *Math. Z.* **195** (1987), 197–204

[8] **Y.T. Siu**—The Fujita conjecture and the extension theorem of Ohsawa-Takegoshi, *Geometric Complex Analysis, Hayama, World Sci. Publ.* (1996)

[9] **Y.T. Siu**—Section extension from hyperbolic geometry of punctured disk and holomorphic families of flat bundles, arxiv:11.042563

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, 35121 PADOVA, ITALY
E-mail address: `baracco@math.unipd.it`